

**ANIMALS, CELLS AND METHODS FOR PRODUCTION OF
DETECTABLY-LABELED ANTIBODIES**

CROSS-REFERENCE TO RELATED APPLICATION

This application claims priority under 35 U.S.C. § 119(e) to provisional application serial no. 60/241,053, filed October 17, 2000, which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The invention is directed to methods for obtaining detectably-labeled antibodies to any antigen by using genetically-modified mammals which express at least one detectably-labeled antibody molecule component.

BACKGROUND OF THE INVENTION

Antibodies are widely used for diagnostic and research purposes for localizing an antigen within a cell, tissue, or other biological sample. For histopathology or cytology, cells are usually first exposed to a primary antibody which is specific for the desired target antigen but not directly detectable. In a subsequent step, the primary antibody is detected with a labeled secondary antibody which recognizes the primary antibody. This procedure is somewhat cumbersome and may result in undesirable background staining due to non-specific reactivity of the secondary antibody. In addition, in order to visualize two or more antigens, the primary antibodies must be from two different species such that each secondary antibody is not cross-reactive; expensive species-specific secondary antibodies must therefore be used for each primary antibody. With

1 advances in optical and digital imaging and detection, multiply-labeled samples and
2 simultaneous detection of several analytes is possible, yet the aforementioned biological
3 limitations do not permit full advantage to be taken of these advances. Moreover, since
4 99monoclonal antibodies are generated typically only in mice and rats, in conventional dual or
5 triple labeling, only one antibody can be monoclonal. This can be an important restriction.

6
7 Several techniques are available to circumvent these limitations, including direct chemical
8 conjugation of the primary antibody with a detectable molecule, or direct conjugation with biotin
9 and subsequent detection with a fluorescent avidin. These procedures are cumbersome and
10 requires purification of the primary antibodies.

11
12 It is towards the facile preparation of a detectable primary antibody to any desired antigen that
13 the present invention is directed.

14
15 The citation of any reference herein should not be construed as an admission that such reference
16 is available as "Prior Art" to the instant application.

17 18 SUMMARY OF THE INVENTION

19 In one broad aspect, the invention is directed to a genetically-modified mammal capable of
20 expressing at least one chimeric immunoglobulin gene comprising at least one detectable protein
21 or peptide fused with a gene expressing an immunoglobulin component selected from the group
22 consisting of the kappa immunoglobulin light chain, the lambda immunoglobulin light chain, an

immunoglobulin heavy chain, and any combination thereof, wherein antibodies secreted by immune cells of the genetically-modified mammal comprise said at least one detectable protein or peptide. The immunoglobulin heavy chain gene may be IgG, IgM, IgD or IgA. In one embodiment, the at least one detectable peptide or protein is present at the C-terminus of the gene product of the fusion polynucleotide; preferably, the at least one detectable peptide or protein present at the C-terminus of the gene product of the fusion polynucleotide is located in exon G1. In another embodiment, the at least one detectable peptide or protein is present at the C-terminus of the gene product of the fusion polynucleotide with a flexible linker peptide therebetween. Preferably, the least one detectable peptide or protein present at the C-terminus of the gene product of said fusion polynucleotide with a flexible linker therebetween is located in exon G1.

In one embodiment, the immunoglobulin molecule secreted by immune cells of the above-mentioned genetically-modified mammal comprises at least one detectable protein or peptide in the heavy chain of the immunoglobulin molecule. In another embodiment, the immunoglobulin molecule secreted by immune cells of the genetically-modified mammal comprises at least one detectable protein or peptide in the light chain of said immunoglobulin molecule. In a further embodiment, the immunoglobulin molecule secreted by immune cells of the genetically-modified mammal comprises at least one detectable protein or peptide in the heavy chain and at least one detectable protein or peptide in the light chain of said immunoglobulin molecule.

In one embodiment of the invention, at least one of the aforementioned detectable proteins or

polypeptides is an autofluorescent protein or peptide, a visibly-detectable protein or peptide, an enzymatically active protein or peptide, a protein or peptide capable of interacting with another molecule to produce a detectable product, or any combination thereof. In another embodiment, the detectable protein or polypeptide is capable of quenching or modulating fluorescence. Non-limiting examples of autofluorescent proteins or peptides include green fluorescent protein, red fluorescent protein, and a fluorescent analog or fragment of any of the foregoing. Green fluorescent protein is preferred. In another embodiment, the at least one detectable protein is a combination of an autofluorescent protein or peptide and an enzymatically-active protein or peptide, such as but not limited to a combination of green fluorescent protein and alkaline phosphatase.

In another broad aspect of the invention, a genetically-modified immune cell is provided which is capable of expressing at least one chimeric immunoglobulin gene comprising at least one detectable protein or peptide fused with a gene expressing an immunoglobulin component selected from the group consisting of the kappa immunoglobulin light chain, the lambda immunoglobulin light chain, an immunoglobulin heavy chain, and any combination thereof, wherein antibodies secreted by the genetically-modified immune cell comprise the at least one detectable protein or peptide. The immunoglobulin heavy chain gene may be IgG, IgM, IgD or IgA. In one non-limiting example, the at least one detectable peptide or protein is present at the C-terminus of the gene product of the fusion polynucleotide; preferably, the polynucleotide encoding the at least one detectable peptide or protein present at the C-terminus of the gene product of said fusion polynucleotide is located in exon G1. In another embodiment, the at least

one detectable peptide or protein present at the C-terminus of the gene product of said fusion polynucleotide has a flexible linker peptide therebetween; preferably, the polynucleotide encoding the at least one detectable peptide or protein present at the C-terminus of the gene product of said fusion polynucleotide with a flexible linker therebetween is located in exon G1.

The immunoglobulin molecule secreted by the aforementioned immune cell may comprise at least one detectable protein or peptide in the heavy chain of the immunoglobulin molecule, or it may comprise at least one detectable protein or peptide in the light chain of said immunoglobulin molecule. In another embodiment, the immunoglobulin molecule secreted by the genetically-modified immune cells described above comprises at least one detectable protein or peptide in the heavy chain and at least one detectable protein or peptide in the light chain of the immunoglobulin molecule.

The aforementioned genetically-modified immune cell may have at least one detectable protein or polypeptide that is capable of quenching or modulating fluorescence; or the protein or peptide is an autofluorescent protein or peptide, a visibly-detectable protein or peptide, an enzymatically active protein or peptide, a protein or peptide capable of interacting with another molecule to produce a detectable product, or any combination thereof. In the example wherein the at least one detectable protein is an autofluorescent protein or peptide, it may be, by way of non-limiting example, green fluorescent protein, red fluorescent protein, or a fluorescent analog or fragment of any of the foregoing. Green fluorescent protein is preferred. In another embodiment, the at least one detectable protein is a combination of an autofluorescent protein or peptide and an

enzymatically-active protein or peptide, such as but not limited to a combination of green fluorescent protein and alkaline phosphatase.

The invention is also directed to a hybridoma comprising the genetically-modified immune cell as mentioned above.

In yet another aspect, the invention is directed to a chimeric, detectably-labeled immunoglobulin molecule comprising at least one detectable protein or peptide fused with the kappa immunoglobulin light chain, the lambda immunoglobulin light chain, an immunoglobulin heavy chain, or any combination thereof. The at least one detectable peptide or protein may be present at the C-terminus of the gene product of said fusion polynucleotide, preferably located in exon G1. A flexible linker peptide may be provided therebetween. In another embodiment, a polynucleotide encoding said at least one detectable peptide or protein present at the C-terminus of the gene product of said fusion polynucleotide with a flexible linker therebetween is located in exon G1. The chimeric, detectably-labeled immunoglobulin molecule may have an immunoglobulin heavy chain gene is selected from IgG, IgM, IgD and IgA. At least one detectable protein or peptide may be present in the heavy chain of said immunoglobulin molecule, the at least one detectable protein or peptide may be present in the light chain of said immunoglobulin molecule, or, in a further embodiment, the at least one detectable protein or peptide may be present in the heavy chain and at least one detectable protein or peptide in the light chain of said immunoglobulin molecule.

1 The aforementioned chimeric, detectably-labeled immunoglobulin molecule may have at least
2 one detectable protein or polypeptide that is capable of quenching or modulating fluorescence;
3 or, the at least one detectable protein or peptide is an autofluorescent protein or peptide, a
4 visibly-detectable protein or peptide, an enzymatically active protein or peptide, a protein or
5 peptide capable of interacting with another molecule to produce a detectable product, or any
6 combination thereof. In one embodiment, the at least one detectable protein is an autofluorescent
7 protein or peptide, such as but not limited to green fluorescent protein, red fluorescent protein, or
8 a fluorescent analog or fragment of any of the foregoing. Preferably, it is green fluorescent
9 protein. In another embodiment, the at least one detectable protein may be a combination of an
10 autofluorescent protein or peptide and an enzymatically-active protein or peptide, such as but not
11 limited to a combination of green fluorescent protein and alkaline phosphatase.

13 In a further broad aspect, the present invention provides a method for producing a quantity of
14 detectably-labelled polyclonal antibodies by carrying out at least the steps of

- 15 a) providing a genetically-modified mammal as described hereinabove;
- 16 b) immunizing the genetically-modified mammal with a preselected immunogen,
17 wherein the genetically-modified mammal generates antibodies to the
18 immunogen, wherein antibodies secreted by immune cells of the genetically-
19 modified mammal comprise the at least one detectable protein or peptide; and
- 20 c) isolating the detectably-labelled antibodies from the genetically-modified
21 mammal.

1 The invention is also directed to a method for producing a quantity of detectably-labelled
2 monoclonal antibodies comprising the steps of

- 3 a) preparing a genetically-modified mammal in accordance with the above
4 description;
- 5 b) immunizing the genetically-modified mammal with a preselected immunogen,
6 wherein immune cells of the genetically-modified mammal generate antibodies to
7 the immunogen, wherein antibodies secreted by the immune cells comprise the at
8 least one detectable protein or peptide;
- 9 c) immortalizing antibody-producing immune cells isolated from the genetically-
10 modified mammal;
- 11 d) selecting immortalized immune cells isolated from the genetically-modified
12 mammal that secrete antibodies specific to the immunogen; and
- 13 e) preparing a quantity of detectably-labeled monoclonal antibodies from the
14 selected immune cells.

15 In a further aspect, the invention is also directed to a genetically-modified mammal capable of
16 producing a detectably-labeled immunoglobulin in response to immunization by an antigen, the
17 genome of the mammal comprising at least one fusion polynucleotide consisting of a
18 polynucleotide sequence encoding at least one detectable protein or peptide fused with a the
19 kappa immunoglobulin light chain gene, the lambda immunoglobulin light chain gene, an
20 immunoglobulin heavy chain gene, or any combination thereof, wherein antibodies secreted by
21 immune cells of the genetically-modified mammal comprise the at least one detectable protein or
22

peptide.

In a further aspect, the invention is directed to a chimeric, detectably-labeled immunoglobulin molecule comprising at least one fluorescent protein or peptide and at least one fluorescence-quenching or -modulating protein or peptide fused with a component of the immunoglobulin molecule independently selected from either the kappa immunoglobulin light chain, the lambda immunoglobulin light chain, an immunoglobulin heavy chain, or any combination thereof.

These and other aspects of the present invention will be better appreciated by reference to the following drawings and Detailed Description.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the beta-galactosidase-binding activity of a chimeric fusion immunoglobulin secreted from 293T cells transfected with nucleic acid encoding a green-fluorescent-protein-labeled heavy immunoglobulin (IgG2) chain and a green-fluorescent-protein-labeled light (kappa) chain hybrid anti-beta-galactosidase immunoglobulin.

Figures 2 A-B show the fluorescence of antibodies in serum from kappa-GFP knock-in mice bound to Protein A/G beads, as compared to those from wild-type mice.

Figure 3 A-B demonstrate using PCR amplification that mice of the invention may be homozygous or heterozygous for the kappa-GFP gene, using kappa light chain primers (Figure

3A). Use of GFP primers (Figure 3B) confirm the presence or absence of the GFP insert.

DETAILED DESCRIPTION OF THE INVENTION

The inventors herein have discovered a novel and facile method for preparing useful quantities of detectably-labeled polyclonal or monoclonal antibodies to any preselected immunogen. The method spares the need to perform any modification of the immunogen-specific (primary) antibody to provide a detectable label thereon, such as conjugation to a fluorophore or enzyme, as the detectably-labeled antibodies of the invention are inherently labeled as they are expressed and secreted from genetically-modified immune system cells. In its broadest aspect, the invention provides genetically-modified animals and immune cells from such animals in which the constant region of the heavy and/or the light chain components of the antibodies produced by the animals or cells is fused with a detectable polypeptide, such as a fluorescent polypeptide, these fusion protein(s) resulting from the expression of a genetic modification of the heavy and/or light chain genes wherein the polynucleotide encoding the detectable label is incorporated into the respective genes. Such genetically-modified animals or cells are provided such that any antibody expressed and secreted by the animal or cell comprises at least one labeled constant region.

Thus, the mere induction of a humoral immune response generates the desired detectably-labeled antibody.

The detectable labels of the invention are polypeptides, such as proteins or peptides, and they may be introduced into the genome of a mammalian organism by such methods as homologous recombination and transfection, but such methods are not intended to be limiting whatsoever, and

1 one of skill in the art may prepare the construct(s) and modified cells in any appropriate manner.

2 Embryonic stem cells may be so modified such that the genome of the animal resulting therefrom
3 comprises the genetic modification. Several examples provided below are for illustrative
4 purposes only.

5
6 The detectable polypeptide may be, by way of non-limiting example, a fluorescent polypeptide, a
7 visibly-colored polypeptide, a polypeptide with enzymatic activity, or a polypeptide capable of
8 interacting with or binding to another molecule to produce a detectable product. Preferred are
9 labels which require no direct interaction or further sample processing for detectability, and thus
10 are detectable by the application of exogenous methods such as absorbance of light, fluorescence,
11 etc. More preferred are fluorescent polypeptides; most preferred is green fluorescent protein and
12 its polypeptide relatives.

13
14 Moreover, a plurality of detectable labels may be present in the antibodies produced by the
15 animals or cells of the invention. For example, both a fluorescent label and an enzyme label may
16 be appended in tandem to the C-terminus of an immunoglobulin heavy chain gene, such that the
17 resulting expressed and assembled immunoglobulin molecule is detectable both by fluorescence
18 and by histochemistry, by use of a fluorogenic and chromogenic/precipitating substrate of the
19 enzyme, respectively. In another example, the antibody is detectable fluorometrically and by
20 Western blot. As noted above, either or both the heavy and light chain constant regions may be
21 independently modified as described herein, both with the same single or plurality of labels, or,
22 for example, the heavy chain with a fluorescent label and the light chain with an enzyme label.

1 These particular examples of multiple labels on multiple sites of the antibodies are merely
2 illustrative of the range of directly-labeled antibodies that the skilled artisan may be directed to
3 prepare following the teachings herein, and any particular example or embodiment is not
4 intended to be limiting whatsoever.

5
6 Examples of fluorescent polypeptides include but are not limited to green fluorescent protein and
7 other related polypeptide fluorophores which may produce other colors; the availability of
8 individually-detectable primary antibodies in a mixture allows for the simultaneous localization
9 or quantitation of multiple target antigens or analytes in a biological sample. The green
10 fluorescent protein of *Aequorea victoria* is particularly preferred as the fluorescent protein. A
11 cDNA for the protein has been cloned (D. C. Prasher et al., "Primary structure of the *Aequorea*
12 *victoria* green-fluorescent protein," *Gene* (1992) 111:229-33.). *Aequorea* green fluorescent
13 protein ("GFP") is a stable, proteolysis-resistant single chain of 238 residues and has two
14 absorption maxima at around 395 and 475 nm; Excitation at the primary absorption peak of 395
15 nm yields an emission maximum at 508 nm with a quantum yield of 0.72-0.85 (O. Shimomura
16 and F. H. Johnson *J. Cell. Comp. Physiol.* 59:223 (1962); J. G. Morin and J. W. Hastings, *J. Cell.*
17 *Physiol.* 77:313 (1971); H. Morise et al. *Biochemistry* 13:2656 (1974); W. W. Ward *Photochem.*
18 *Photobiol. Reviews* (Smith, K. C. ed.) 4:1 (1979); A. B. Cubitt et al. *Trends Biochem. Sci.*
19 20:448-455 (1995); D. C. Prasher *Trends Genet.* 11:320-323 (1995); M. Chalfie *Photochem.*
20 *Photobiol.* 62:651-656 (1995); W. W. Ward. *Bioluminescence and Chemiluminescence* (M. A.
21 DeLuca and W. D. McElroy, eds) Academic Press pp. 235-242 (1981); W. W. Ward & S. H.
22 Bokman *Biochemistry* 21:4535-4540 (1982); W. W. Ward et al. *Photochem. Photobiol.*

35:803-808 (1982)). Mutants of GFP are embraced herein as they provide certain other characteristics, such as mutation of Serine 65 to Thr (S65T) simplifies the excitation spectrum to a single peak at 488 nm of enhanced amplitude (R. Heim et al. Nature 373:664-665 (1995)), which no longer gives signs of conformational isomers (A. B. Cubitt et al. Trends Biochem. Sci. 20:448-455 (1995)). In another example, U.S. Patent No. 6,077,707 describes a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein but differs by at least a substitution at T203 and, in particular, T203X, wherein X is an aromatic amino acid selected from H, Y, W or F. In one embodiment therein, the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I. In another embodiment, the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S65G/V68L/Q69K/S72A/T203Y; S72A/S65G/V68L/T203Y; S65G/S72A/T203Y; or S65G/S72A/T203W. In another embodiment, the nucleotide sequence encoding the protein differs from the nucleotide sequence of native green fluorescent protein by the substitution of at least one codon encoding an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222 (not E222G), or V224. In another embodiment, the amino acid substitution is: L42X, wherein X is selected from C, F, H, W and Y, V61 X, wherein X is selected from F, Y, H and C, T62X, wherein X is selected from A, V, F, S, D, N, Q, Y, H and C, V68X, wherein X is selected from F, Y and H, Q69X, wherein X is selected from K, R, E and G, Q94X, wherein X is selected from D, E, H, K and N, N121X, wherein X is selected from F, H, W and Y, Y145X, wherein X is selected from

W, C, F, L, E, H, K and Q, H148X, wherein X is selected from F, Y, N, K, Q and R, V150X, wherein X is selected from F, Y and H, F165X, wherein X is selected from H, Q, W and Y, I167X, wherein X is selected from F, Y and H, Q183X, wherein X is selected from H, Y, E and K, N185X, wherein X is selected from D, E, H, K and Q, L220X, wherein X is selected from H, N, Q and T, E222X, wherein X is selected from N and Q, or V224X, wherein X is selected from H, N, Q, T, F, W and Y. These examples are merely illustrative of the wide selection of fluorescent polypeptides and their corresponding polynucleotide sequences that may be employed in the preparation of the genetically-modified cells or animals of the invention.

Another example of a fluorescent protein is red fluorescent protein from coral (Matz et. al., 1999, *Nature Biotechnology* 17:969). Fluorescent peptides or fluorescent protein fragments of these and other fluorescent proteins are also embraced herein. In another example of a detectable polypeptide, polypeptide labels capable of being detected include those containing four cysteines at the i, i+1, i+4, and i+5 positions (i.e., WEAAAREACCRECCARA). These peptides will bind specifically to the fluorescein derivative 4',5'-bis(1,3,2-dithioarsolan-2-yl)fluorescein (FLASH), which is non-fluorescent until bound. The use of such peptides has the advantage of providing fusion proteins of much smaller size than GFP, and fluorescence is easily detected by addition of FLASH (Griffin et. al., 1998, *Science* 281:269). Thus, the use of polypeptides that are capable of specifically binding a detectable label are embraced herein as another embodiment of the present invention.

Examples of polypeptides capable of being detected by enzymatic activity include various

1 enzymes and catalytic polypeptide fragments thereof. Particularly preferred enzymatic labels
2 include those which are able to produce a detectable color or fluorophore in a single step
3 requiring a minimum of reagents, such as alkaline phosphatase, which can cleave a chromogenic
4 substrate, such as p-nitrophenyl phosphate, or a fluorogenic substrate, such as ECF substrate
5 (Amersham/Pharmacia); and a fluorogenic horseradish peroxidase substrate, FluoroBlot (Pierce
6 Chemical Co.). Another example is a fluorescent beta-galactosidase substrate that can be used in
7 live cells and is 100-fold more sensitive than GFP (Zlokarnik et. al., 1988, *Science* **279**:85). The
8 skilled artisan by the teachings herein will be amply aware of polynucleotides that may be fused
9 to the immunoglobulin constant region(s) and upon expression produce an enzymatically-active
10 fusion polypeptide comprising the immunoglobulin heavy and/or light chain.

11 Such labels may also result in the precipitation of a substrate, for histochemical localization of
12 the antibody, or the label may interact with another detectable component, and thus be, for
13 example, an intein, a biotin-binding subunit of streptavidin or avidin, a His tag, or a chitin-
14 binding domain.
15

16
17 Guidance for the selection of the immunoglobulin genes and locations therein in which to fuse
18 the detectable polypeptide(s) of the invention may be performed as follows. As is well known,
19 the antibody molecule is composed of two heavy chains and two light chains. Each chain has a
20 two identical variable regions which bind antigen, and a constant region. The N-terminus of both
21 chains is part of the variable region and the C-terminus is part of the constant region.
22

1 The genomic organization of heavy chain contains a number of repeats of V, D, and J regions
2 upstream of several constant regions. The different constant regions define the eventual antibody
3 class (e.g., IgG1, IgG2a, IgA, etc). During B-cell maturation, a single V, D, and J are joined to
4 form a single exon. Additionally, late in B-cell development, a single constant region is chosen in
5 a process called "class switching". Each constant region contains a number of exons. After
6 transcription, the VDJ and constant regions are spliced into the mature mRNA.

7
8 To generate the fusion protein of the invention, a polynucleotide encoding the detectable
9 polypeptide is inserted into the constant region, or preferably, is appended to portion of the gene
10 resulting in the expression of the fused polypeptide label on the C-terminus of the constant region
11 of the light and/or heavy chains, optionally with a flexible linker peptide.

12
13 There are two classes of light chains, kappa and lambda. Kappa comprises 95% of antibodies,
14 lambda 5%. Furthermore, there are two forms of each class of the heavy chain—a secreted and a
15 membrane bound form. The membrane bound form is responsible for signaling to the B-cell
16 when it is crosslinked by an antigen and is required for B-cell maturation and antibody
17 production. The two forms result from alternative splicing. Preferably, only the secreted form is
18 modified as described herein. The membrane bound form would be completely wild type, to
19 maintain fidelity of generation of the antibody response.

20
21 Thus, an immune cell with the genetic modifications described herein may secrete
22 immunoglobulin that may have possess a detectable polypeptide label fused in the heavy chain

constant region, or a detectable polypeptide label fused in the light chain constant region, or the immunoglobulin may have both a heavy and light chain fusion component. Moreover, each detectable label may be a single detectable label or a plurality of labels, in tandem or not, optionally separated from the immunoglobulin portion of the polypeptide by one or more linker sequences. The label on the heavy chain may be the same or different from that on the light chain, for immunoglobulins that have both heavy and light chains as fusion products. Although the placement of the detectable label is preferably on or near the C-terminus, the fused label(s) may be at any position(s) with does not detract from the ability of the immunoglobulin to interact with and bind to its target antigen. The invention herein embraces any and all variations in the position, placement, linkers, and number of fused peptides or polypeptides, with the object of the invention to provide at least one detectable label on an immunoglobulin molecule generated by exposure of an animal or immune cell with the aforementioned genetic modification to any immunogen capable of eliciting secretion of an antibody directed thereto.

As described above, a mammalian organism, such as a mouse, rat, rabbit, goat, cow, horse, may be provided with the genetic modification as described herein. Example 4 below demonstrates the invention using mice, in which endogenous production of fluorescent antibodies is shown. Exposure of the animal to a preselected immunogen will elicit an antibody response, the secreted antibodies having the detectable label(s) present in every antibody molecule. For production of polyclonal antibodies, antiserum from the immunized animals may be collected, the antibodies purified if desired, for subsequent use in such areas as histochemistry, diagnostics, etc.

1 If detectably-labeled primary monoclonal antibodies are desired, routine methods may be
2 followed using an animal, such as a mouse, with the genetic makeup hereindescribed. B-cells
3 secreting the detectably-labeled primary antibodies may be fused for immortalization, and
4 screening and selection for stable antibody-secreting hybridomas obtained by routine methods.

5
6 Of course, detectably-labeled primary anti-idiotypic antibodies may be prepared by the foregoing
7 methods, using antibodies as immunogen.

8
9 As mentioned above, the polyclonal and monoclonal antibodies generated by the methods
10 described herein have several advantages, including 1) because the primary antibody is
11 endogenously fluorescent, no secondary antibody or chemical labeling is required, 2)
12 autofluorescent proteins are currently available in several different colors, including blue, cyan,
13 green, yellow, and red, allowing for simultaneous labeling of several antigens, and simultaneous
14 detection by instrumentation capable of discriminating several fluorophores simultaneously. In
15 addition, the availability of two different means for detection, such as but not limited to a visibly-
16 detectable and enzymatically-detectable marker, provides detectability under a variety of
17 conditions, applicable to various research and diagnostic applications, among others.

18
19 Introduction of the genetic modification(s) described herein into an embryonic stem cell or other
20 cell type may be achieved by a variety of procedures known in the art. Preferably, the genetic
21 modification is introduced by a "knock-in" procedure in which the wild-type immunoglobulin
22 gene(s) are replaced by the detectable-label-modified genes. Such procedures may include the

1 use of a bacterial artificial chromosome, as exemplified in Example 2 below, although this
2 example is merely illustrative and non limiting as to procedures for achieving the genetically-
3 modified mammal or mammalian cells of the invention.
4

5 Antigens to which detectable antibodies may be raised in genetically-modified animals or
6 immune cells as described herein are not limited to any particular types or classes of immunogen,
7 and includes those for which a detectable antibody is desirable. Such antigens comprise a vast
8 list. By way of non-limiting example, this includes diagnostic and research reagents for
9 identifying the presence of and/or quantitating various medically-important biomolecules in
10 bodily fluids, biopsy and necropsy samples, for example, all diagnostic tests which employ
11 immunoassay protocols, including ELISA, radioimmunoassay, EMIT, immunofluorescence,
12 fluorescence polarization, and other methods for detecting the interaction between and
13 biomolecule of interest and an antibody thereto. Medical diagnostics include, by way of non-
14 limiting example, assays for autoimmune disorders, cardiovascular disorders, diabetes, endocrine
15 disorders, fungal, bacterial, viral, parasitic, and other infectious agents, hematologic diseases,
16 immunologic diseases, hepatic diseases, oncologic diseases, thyroid diseases, and toxicology and
17 drugs of use and abuse. Moreover, as mentioned above, the ability to discriminate between
18 differently labeled primary antibodies without the need for, and concomitant disintegration of
19 signal, by multiple secondary antibodies, will permit multiple analytes to be measured
20 simultaneously, decreasing the costs and increasing the amount of information available for
21 rendering diagnostic and therapeutic decisions. In the research area, innumerable new and
22 known biomolecules to which antibodies are routinely raised for identifying the location,

1 movement, transport, role, etc., of such new or rediscovered biomolecules in biological processes
2 will be simplifiable, as well as permit the simultaneous and facile monitoring of multiple
3 biomolecules using differently-labeled antibodies to several biomolecules participating in a
4 process. The discussion herein on particular antigens is not intended to be at all limiting but is
5 merely illustrative of some examples of the utility of the invention.

6
7 In another example of the invention, a genetically-modified mammal may be prepared which
8 responds to immunization with an antigen by the production of chimeric immunoglobulin
9 molecules capable of reading out a signal or altering the signal produced only on binding with the
10 target antigen. This signal may be, for example, an increase, decrease or change in fluorescence.
11 Such chimeric antibodies may be used in a very simple homogeneous immunoassay in which, on
12 combining with a sample, indicates the presence or extent of the level of the antigen in the
13 sample. Pairs of proteins or peptides capable of undergoing such modulation in fluorescence
14 include FRET pairs, such as described, for example, in U.S. Patent 5,998,204, incorporated
15 herein by reference in its entirety.
16

17 Such chimeric antibodies are prepared by following the methods described herein. In the
18 example wherein the fluorescence is detected, the genes encoding the chains of the secreted
19 immunoglobulin are modified at the polynucleotide level to provide that both a fluorescent
20 peptide or polypeptide, and a fluorescence-quenching or -modulating peptide or polypeptide, are
21 fused into either the heavy immunoglobulin chain or the light immunoglobulin chain in the
22 appropriate position in the genome of the mouse. The positions of integration into the respective

immunoglobulin components are such that the proximity of the fluorescent polypeptide and the fluorescence-quenching or -modulating polypeptide in the secreted, whole immunoglobulin molecule, change on binding of the immunoglobulin to its target antigen. This change in proximity alters the interaction between the fluorophore and the quencher or modulator resulting in a modulation in the detectable fluorescence on exposure of the immunoglobulin to its excitation wavelength. Such positions are selected to not alter the ability of the complete immunoglobulin molecule to assemble, be secreted, or bind the antigen. These selections are within the realm of the skilled artisan. Known fluorescent peptides or polypeptides as well as peptides and polypeptides capable of quenching or modulating the fluorescence, such as FRET pairs, are known and can be selected to provide the chimeric immunoglobulin, and mammals capable of producing the immunoglobulin after immunization.

The present invention may be better understood by reference to the following non-limiting Examples, which are provided as exemplary of the invention. The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as limiting the broad scope of the invention.

Example 1

Secretion of detectably-labeled antibodies

A rat IgG2A antibody hybridoma which produces a monoclonal antibody reactive against beta-galactosidase was used to determine whether GFP fusion proteins would be correctly

1 assembled, secreted, and recognize antigen. The cDNA of the secreted form of the heavy and
2 light chains against beta-galactosidase were cloned. The polynucleotide sequence encoding GFP
3 was placed in front of the stop codon of both the heavy chain of IgG2A and light kappa chains.
4 The plasmid combinations of 1) non-GFP tagged, 2) IgG2A heavy chain tagged, 3) light chain
5 tagged, and 4) both heavy chain and light chain tagged, were transfected into 293T kidney cells
6 and tested for ability to bind antigen.

7
8 The 293T cells do not express any endogenous antibodies. They were transfected with two
9 expression vectors of combinations of IgG2A and kappa chains against beta-galactosidase: 1)
10 IgG2A and kappa; 2) IgG2A and kappa-GFP; 3) IgG2A-GFP and kappa; and 4) IgG2A-GFP and
11 kappa-GFP. The transfected cells will express and assemble and secrete the complete antibody.
12 Forty-eight hours after transfection, the medium was harvested. This contained unpurified
13 antibodies which was used for the experiments below

14
15 Confirmation of the secretion of the detectably-labeled antibody was obtained in several
16 experiments. In one experiment, cells were fed radiolabeled ^{35}S methionine after transfection and
17 newly synthesized proteins were labeled. The secreted proteins were collected. Two fractions
18 were evaluated, one that bound to Protein G and one that bound to beta-galactosidase
19 immobilized on beads. Protein G is a toxin which binds antibodies. As shown in **Figure 1**, both
20 GFP-tagged and untagged antibodies bound to Protein G and to beta-galactosidase. The affinity
21 and specificity towards beta-galactosidase appears greater. Thus, the antibody recognizes antigen
22 and is correctly folded. When only light chain or heavy chain but not both are transfected, the

1 chains did not bind to either protein G or beta-galactosidase. When examined under the
2 microscope, the beads were bright green, showing that the GFP is correctly folded and
3 fluorescent.

4 **Example 2**

5 **Preparation of genetically-modified (knock-in) animals comprising** 6 **detectably-labeled immunoglobulin genes**

7
8 Knock-in or gene replacement involves replacing an endogenous piece of DNA in the
9 chromosome with a constructed piece of DNA. Thus, the constructed DNA must correctly
10 integrate into the same genetic location as the gene to be replaced. This is distinctly different
11 from transgenic technology where a constructed DNA is randomly introduced into a cell.

12
13 I. Construction of a segment of bacterial plasmid DNA that contains the change of interest and
14 extensive homology to the target. Generation of the IgG1-GFP fusion knock-in construct is
15 described below. The VDJ variable regions of the IgG1 constant chain are far upstream (5') of
16 this area. Immediately upstream of this area are the IgM and IgD constant regions. Downstream
17 to this area are the constant regions for IgG2a, IgG2b, IgG3, IgE, and IgA. The exons CH1, CH2
18 and G1 are spliced together to make the constant region of the secreted form. To make the
19 membrane bound form, CH1, CH2, G1 (except the last 2 amino acids), TD1, and TD2 are spliced
20 together. In the targeting vector, GFP is fused in-frame at the end of the G1 exon with a five-
21 amino-acid linker of Gly-Gly-Ser-Gly-Gly in-between. The membrane bound form splices out
22 the last two amino acids of G1 and thus the entire GFP. Therefore, GFP will be fused only to the

1 secreted form.

2
3 A bacterial artificial chromosome (BAC) containing the entire IgG region is obtained. The
4 targeting vector contains the 5' homologous region directly PCR'd from the BAC. The G1-GFP
5 fusion is generated by primer overlap extension PCR (Horton et al., 1990; Horton, 1995). A
6 neomycin resistance gene is introduced as a selection marker. This neomycin gene is flanked on
7 each side by a loxP site. Thus, after selection, the gene is deleted by Cre-mediated
8 recombination. This eliminates problems due to the presence of the neomycin gene and its
9 associated promoter (Zou et al., 1993). A 3' homologous region is also PCR'd from the BAC.
10 Finally, a toxic gene, diphtheria toxin (DTA) is introduced just outside the area of homology.
11 Homologous recombination between the targeting vector and the targeting region cuts out the
12 toxin gene. Non-homologous random integration events often retain DTA and the cells will be
13 killed (Yagi et al., 1990; Yagi et al., 1993; McCarrick, III et al., 1993).

14
15 After preparation, embryonic stem cells are transfected with the targeting vector and selection
16 with neomycin analog G418 is performed. Clones are screened for correct integration by
17 Southern analysis and PCR. Subsequently, correctly targeted ES cells are injected into blastocysts
18 and implanted into mothers. The blastocysts develop into chimeric mice, where some cells are
19 developed from the injected ES cells. The chimeras are used to parent heterozygote mice. Two
20 heterozygous mice are then parented to produce homozygous mice.

21
22 The mouse IgG1 heavy chain and kappa light chains have been previously targeted for gene

1 replacement. In those cases, the constant chain was replaced with human constant chains and the
2 resulting mice generated humanized antibodies (Zou et al., 1993; Zou et al., 1994).

3
4 **Production of Monoclonal Antibodies.** Mice as prepared above are inoculated with the antigen
5 of interest. After several weeks, the antibody production against the antigen is screened, as
6 traditionally done with an ELISA, where the antigen is immobilized, the mouse serum is added,
7 and an enzyme-linked secondary antibody against the primary is added. The presence of
8 enzymatic activity indicates that the mouse has produced specific antibodies. Since here, the
9 antibodies are already fluorescent, an ELISA is unnecessary and measurement of bound
10 fluorescence is sufficient. Fusion of splenocytes from a positive mouse with myeloma cells is
11 used to produce immortalized antibody-producing cells, following standard protocols. Further
12 screening of colonies can be performed using simple fluorescence measurements.

13
14 Antibody may be produced in quantity by growth of cell-hybrids, following standard protocols.

15 **Example 3**

16 **Homogeneous immunoassays using detectably-labeled antibodies**

17 **from a genetically-modified mouse**

18
19
20 In a further example of the methods described in Example 2, above, a genetically-modified
21 mouse may be prepared which responds to immunization with an antigen by the production of
22 chimeric immunoglobulin molecules capable of reading out a signal only on binding with the

1 target antigen. Such chimeric antibodies may be used in a very simple homogeneous
2 immunoassay in which, on combining with a sample, indicates the presence or extent of the level
3 of the antigen in the sample.

4 Such chimeric antibodies are prepared by following the methods herein. Both a fluorescent
5 peptide or polypeptide, and a fluorescence-quenching peptide or polypeptide, are fused into
6 either the heavy immunoglobulin chain or the light immunoglobulin chain in the appropriate
7 position in the genome of the mouse. Such pairs may include FRET pairs or proteins, such as
8 described, for example, in U.S. Patent 5,998,204, incorporated herein by reference in its entirety.

9 The positions of integration into the respective immunoglobulin components are such that the
10 proximity of the fluorescent polypeptide and the fluorescence-quenching polypeptide change on
11 binding of the immunoglobulin to its target antigen. Such positions also do not alter the ability
12 of the complete immunoglobulin molecule to assemble, be secreted, or bind the antigen. Both
13 the fluorescent and the fluorescence-quenching protein or peptides may be fused into the same
14 immunoglobulin component, at positions wherein antigen binding induces a conformational
15 change and thus a proximity change among the pair and attendant modulation of fluorescence.
16

17 Examples of fluorescent peptides and proteins are described above. In this example, the mouse
18 was genetically modified such that IgG2 heavy chains included the). Suitable pairs, for example
19 include a blue-shifted GFP mutant P4-3 (Y66H/Y145F) as the donor, and an improved green
20 mutant S65T can respectively serve as a donor and an acceptor for fluorescence resonance energy
21 transfer (FRET; Tsien et al., 1993, Trends Cell Biol. 3:242-245). A genetically-modified mouse
22 expressing antibodies with these modifications was immunized with human chorionic

gonadotropin (hCG). The fluorescence of antibodies produced by this mouse was increased upon binding to hCG. This reagent was used in a simplified assay in an automated fluorescence-based instrument for determining pregnancy among a large battery of other diagnostic tests on blood and urine samples.

Example 4

Kappa-GFP knock-in mice produce fluorescent endogenous antibodies

Kappa-GFP knock-in mice were prepared in accordance with the methods described in Example 2, above. At 8 weeks of age, animals were bled and serum was incubated with Protein A/G beads, to which antibodies in the serum bind. As shown in **Figure 2**, under fluorescence illumination the beads showed the characteristic fluorescence of GFP (**Figure 2A**), whereas serum from wild-type mice incubated with Protein A/G beads did not (**Figure 2B**). Thus, the mice produced fluorescently-labeled, endogenous immunoglobulin molecules, comprising kappa-GFP, fluorescent light chains.

Example 5

Confirmation of genotype using PCR for kappa and GFP

Figure 3 shows the results of PCR amplification using primers from up- and down-stream of the section of the kappa light chain where the GFP was targeted (Figure 3A), or using GFP primers (Figure 3B), from DNA extracts from a number of mice prepared in accordance with the present

invention. The segment amplified from wild type kappa light chains is the smallest and runs the fastest on the gel (as indicated). The segment amplified with the GFP insert is larger and migrates slower.

In Figure 3A, using the kappa primers, lane 2 has only the faster moving product and is therefore from a mouse that is homozygous for wild-type kappa light chain. Lanes 3, 5, and 7-12 have both products, which indicates that the mice are heterozygous. Lanes 1, 4, and 6 only have the slower moving product indicating that they are homozygous for the kappa with the inserted GFP.

Figure 3B shows using primers from the GFP coding region that lane 2 was negative for GFP, consistent with the observation that this mouse did not produce a PCR product from kappa-GFP. Lanes 1 and 2-12 were positive for GFP, consistent with the results from the top panel.

The present invention is not to be limited in scope by the specific embodiments describe herein.

Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and the accompanying figures.

Such modifications are intended to fall within the scope of the appended claims.

It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

Horton,R.M. (1995). PCR-mediated recombination and mutagenesis. SOEing together tailor-made genes. Mol. Biotechnol. 3, 93-99.

Horton,R.M., Cai,Z.L., Ho,S.N., and Pease,L.R. (1990). Gene splicing by overlap extension: tailor-made genes using the polymerase chain reaction. Biotechniques 8, 528-535.

McCarrick,J.W., III, Parnes,J.R., Seong,R.H., Solter,D., and Knowles,B.B. (1993). Positive-negative selection gene targeting with the diphtheria toxin A- chain gene in mouse embryonic stem cells. Transgenic Res. 2, 183-190.

Yagi,T., Ikawa,Y., Yoshida,K., Shigetani,Y., Takeda,N., Mabuchi,I., Yamamoto,T., and Aizawa,S. (1990). Homologous recombination at c-fyn locus of mouse embryonic stem cells with use of diphtheria toxin A-fragment gene in negative selection. Proc. Natl. Acad. Sci. U. S. A 87, 9918-9922.

Yagi,T., Nada,S., Watanabe,N., Tamemoto,H., Kohmura,N., Ikawa,Y., and Aizawa,S. (1993). A novel negative selection for homologous recombinants using diphtheria toxin A fragment gene. Anal. Biochem. 214, 77-86.

1 Zou,Y.R., Gu,H., and Rajewsky,K. (1993). Generation of a mouse strain that produces
2 immunoglobulin kappa chains with human constant regions [see comments]. Science 262,
3 1271-1274.

4
5 Zou,Y.R., Muller,W., Gu,H., and Rajewsky,K. (1994). Cre-loxP-mediated gene replacement: a
6 mouse strain producing humanized antibodies. Curr. Biol. 4, 1099-1103.

7

8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211